

Smart Phrase for prenatal aspirin screening used at Boston Medical Center (.BMCOBPNA)

SmartPhrase Editor ? ✕

Name: ID: 222377 Type: User

Content | Owners & Users | Synonyms Rich text (bold, italics, etc.)

Do not include PHI or patient-specific data in SmartPhrases.

1 2 3 4 5 6 7 8 9

Patient was screened for risk of preeclampsia.

At least **1** Major Risk Factor: {AM - EX Major Risk Factors:21148} ←

Several Moderate Risk Factors: {EX Moderate Risk Factors:21157} ←

Patient Qualifies for prenatal aspirin? {Qualify for PNA?:21159} ←

Patient Prescribed prenatal aspirin? {Qualify for PNA?:21159} ←

In EPIC, creates a drop down of selectable criteria based on USPSTF risk factors

Insert SmartList:

Connection logic in this SmartPhrase:

Show label in this SmartPhrase:

Short Description
(250 characters max.)

[Populate from SmartPhrase text](#)

Patient was screened for risk of preeclampsia.

At least 1 Major Risk Factor: {AM - EX Major Risk Factors:21148}

Several Moderate Risk Factors: {EX Moderate Risk Factors:21157}

Patient Qualifies for prenatal aspirin? {Qualify for PNA?:21159}

Embedded SmartLink Format

?

Smart Phrase for counseling postpartum patients for prenatal aspirin use in the next pregnancy (.OBASPIRINNEXTPREGNANCY)

SmartPhrase Editor ? ✕

Name: ID: 210019 Type: User

Content Owners & Users Synonyms Rich text (bold, italics, etc.)

Do not include PHI or patient-specific data in SmartPhrases.

1 2 3 4 5 6 7 8 9

As current pregnancy, labor, or post partum period was complicated by preeclampsia, the patient was counseled about her increased risk for complications of high blood pressure in a future pregnancy. We discussed the need for prenatal aspirin (81 mg) beginning at 12 weeks in a future pregnancy. Discussed how prenatal aspirin has a significant risk reduction of preeclampsia, preterm labor, and IUGR.

Insert SmartList:

Connection logic in this SmartPhrase:

Show label in this SmartPhrase:

+ Add to SmartPhrase

Short Description
(250 characters max.)

[Populate from SmartPhrase text](#)

As current pregnancy, labor, or post partum period was complicated by preeclampsia, the patient was counseled about her increased risk for complications of high blood pressure in a future pregnancy. We discussed the need for prenatal aspirin (81 m...

Embedded SmartLink Format

?